



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C07D 215/56, 417/06, A61K 31/47, A61P 31/22</p>	A1	<p>(11) International Publication Number: WO 00/40563</p> <p>(43) International Publication Date: 13 July 2000 (13.07.00)</p>											
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top; padding: 5px;"> <p>(21) International Application Number: PCT/US99/27959</p> <p>(22) International Filing Date: 22 December 1999 (22.12.99)</p> <p>(30) Priority Data:</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 30%;">60/115,113</td> <td style="width: 40%;">8 January 1999 (08.01.99)</td> <td style="width: 30%;">US</td> </tr> <tr> <td>60/138,390</td> <td>9 June 1999 (09.06.99)</td> <td>US</td> </tr> <tr> <td>60/140,614</td> <td>23 June 1999 (23.06.99)</td> <td>US</td> </tr> </table> <p>(71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): TURNER, Steven, Ronald [US/US]; 1713 Banbury Road, Kalamazoo, MI 49001 (US). STROHBACH, Joseph, Walter [US/US]; 54490 M-66, Mendon, MI 49072 (US). THAISRIVONGS, Suvit [US/US]; 5695 Swallow, Kalamazoo, MI 49002 (US). VAILLANCOURT, Valerie, A. [US/US]; 4342 Bronson Boulevard, Kalamazoo, MI 49008 (US). SCHNUTE, Mark, E. [US/US]; 4459 Wimbledon Way, Kalamazoo, MI 49009 (US). SCOTT, Allen [US/US]; 5287 Saddle Club Drive, Kalamazoo, MI 49009 (US).</p> </td> <td style="width: 50%; vertical-align: top; padding: 5px;"> <p>(74) Agent: YANG, Lucy, X.; Intellectual Property Legal Services, Pharmacia & Upjohn Company, 301 Henrietta Street, Kalamazoo, MI 49001 (US).</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> </td> </tr> </table>			<p>(21) International Application Number: PCT/US99/27959</p> <p>(22) International Filing Date: 22 December 1999 (22.12.99)</p> <p>(30) Priority Data:</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 30%;">60/115,113</td> <td style="width: 40%;">8 January 1999 (08.01.99)</td> <td style="width: 30%;">US</td> </tr> <tr> <td>60/138,390</td> <td>9 June 1999 (09.06.99)</td> <td>US</td> </tr> <tr> <td>60/140,614</td> <td>23 June 1999 (23.06.99)</td> <td>US</td> </tr> </table> <p>(71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): TURNER, Steven, Ronald [US/US]; 1713 Banbury Road, Kalamazoo, MI 49001 (US). STROHBACH, Joseph, Walter [US/US]; 54490 M-66, Mendon, MI 49072 (US). THAISRIVONGS, Suvit [US/US]; 5695 Swallow, Kalamazoo, MI 49002 (US). VAILLANCOURT, Valerie, A. [US/US]; 4342 Bronson Boulevard, Kalamazoo, MI 49008 (US). SCHNUTE, Mark, E. [US/US]; 4459 Wimbledon Way, Kalamazoo, MI 49009 (US). SCOTT, Allen [US/US]; 5287 Saddle Club Drive, Kalamazoo, MI 49009 (US).</p>	60/115,113	8 January 1999 (08.01.99)	US	60/138,390	9 June 1999 (09.06.99)	US	60/140,614	23 June 1999 (23.06.99)	US	<p>(74) Agent: YANG, Lucy, X.; Intellectual Property Legal Services, Pharmacia & Upjohn Company, 301 Henrietta Street, Kalamazoo, MI 49001 (US).</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p>
<p>(21) International Application Number: PCT/US99/27959</p> <p>(22) International Filing Date: 22 December 1999 (22.12.99)</p> <p>(30) Priority Data:</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 30%;">60/115,113</td> <td style="width: 40%;">8 January 1999 (08.01.99)</td> <td style="width: 30%;">US</td> </tr> <tr> <td>60/138,390</td> <td>9 June 1999 (09.06.99)</td> <td>US</td> </tr> <tr> <td>60/140,614</td> <td>23 June 1999 (23.06.99)</td> <td>US</td> </tr> </table> <p>(71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): TURNER, Steven, Ronald [US/US]; 1713 Banbury Road, Kalamazoo, MI 49001 (US). STROHBACH, Joseph, Walter [US/US]; 54490 M-66, Mendon, MI 49072 (US). THAISRIVONGS, Suvit [US/US]; 5695 Swallow, Kalamazoo, MI 49002 (US). VAILLANCOURT, Valerie, A. [US/US]; 4342 Bronson Boulevard, Kalamazoo, MI 49008 (US). SCHNUTE, Mark, E. [US/US]; 4459 Wimbledon Way, Kalamazoo, MI 49009 (US). SCOTT, Allen [US/US]; 5287 Saddle Club Drive, Kalamazoo, MI 49009 (US).</p>	60/115,113	8 January 1999 (08.01.99)	US	60/138,390	9 June 1999 (09.06.99)	US	60/140,614	23 June 1999 (23.06.99)	US	<p>(74) Agent: YANG, Lucy, X.; Intellectual Property Legal Services, Pharmacia & Upjohn Company, 301 Henrietta Street, Kalamazoo, MI 49001 (US).</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p>			
60/115,113	8 January 1999 (08.01.99)	US											
60/138,390	9 June 1999 (09.06.99)	US											
60/140,614	23 June 1999 (23.06.99)	US											
<p>(54) Title: 4-OXO-1,4-DIHYDRO-3-QUINOLINECARBOXAMIDES AS ANTIVIRAL AGENTS</p>													
<p>(57) Abstract</p> <p>The present invention provides a compound of formula (I) Wherein R¹ is C₁₋₇ alkyl, optionally substituted by hydroxy or NR⁴R⁵; R² is C₁₋₇ alkyl substituted by hydroxy or NR⁴R⁵; R³ is H, F or C₁₋₇ alkoxy; R⁴ and R⁵ together with N are a 5- or 6-membered heterocyclic moiety having 1-3 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur in which sulfur may be substituted by one (1) or two (2) oxygen atoms; and pharmaceutically acceptable salts thereof. Compounds of formula (I) of the present invention are useful as antiviral agents.</p>													
<div style="position: absolute; right: 0; top: 50%; transform: translateY(-50%); font-weight: bold;">(I)</div>													

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

4-OXO-1,4-DIHYDRO-3-QUINOLINECARBOXAMIDES AS ANTIVIRAL AGENTS

FIELD OF THE INVENTION

The present invention provides 4-oxo-1,4-dihydro-3-quinolinecarboxamide
5 derivatives, more specifically, provides (4-chlorobenzyl)-4-oxo-1,4-dihydro-3-
quinolinecarboxamide of formula I. These compounds are useful as antiviral agents, in
particular, as agents against viruses of the herpes family.

BACKGROUND OF THE INVENTION


10 The herpesviruses comprise a large family of double stranded DNA viruses.
They are the source of the most common viral illnesses in man. Eight of the
herpesviruses, herpes simplex virus types 1 and 2 (HSV-1 and HSV-2), varicella zoster
virus (VZV), human cytomegalovirus (HCMV), Epstein-Barr virus (EBV), and human
herpes viruses 6, 7, and 8 (HHV-6, HHV-7, and HHV-8), have been shown to infect
15 humans.

HSV-1 and HSV-2 cause herpetic lesions on the lips and genitals, respectively.
They also occasionally cause infections of the eye and encephalitis. HCMV causes birth
defects in infants and a variety of diseases in immunocompromised patients such as
retinitis, pneumonia, and gastrointestinal disease. VZV is the causative agent of chicken
20 pox and shingles. EBV causes infectious mononucleosis. It can also cause lymphomas
in immunocompromised patients and has been associated with Burkitt's lymphoma,
nasopharyngeal carcinoma, and Hodgkins disease. HHV-6 is the causative agent of
roseola and may be associated with multiple sclerosis and chronic fatigue syndrome.
HHV-7 disease association is unclear, but it may be involved in some cases of roseola.
25 HHV-8 has been associated with Kaposi's sarcoma, body cavity based lymphomas, and
multiple myeloma.

Compounds of the present invention are distinct from all other hydroxyquinoline
pharmaceutical agents in that the unique position of chloro substituent on the N-
phenylmethyl of formula I is critical for having useful antiviral activities. These
30 compounds are useful to treat or prevent the above referenced herpesviral infections,
particularly, human cytomegaloviral infection.

INFORMATION DISCLOSURE


US patent 5,891,878 discloses the use of compounds of structure 1 for the treatment of a disease state capable of being modulated by inhibition of production of phosphodiesterase IV or tumor necrosis factor,



5

wherein R^1 may be alkyl(1-6), alkyl(1-6)cycloalkyl, alkyl(1-6)heterocycle, alkyl(1-6)aryl, or alkyl(1-6)heteroaryl each optionally substituted by halogen, alkoxy, hydroxy, CN, CO_2H or corresponding ester or amide, alkyl(1-6), NR^9R^{10} , SO_2NR_2 ; R^3 may be phenyl, pyridyl, furyl, pyrazinyl, pyridazinyl, pyrimidinyl, or cycloalkyl(3-10) which may be
 10 optionally substituted by halogen, alkoxy(1-6), OH, CN, CO_2H and corresponding esters or amides, alkyl(1-6), haloalkyl(1-6), NR^9R^{10} , SO_2NR_2 , aryl, heteroaryl, cycloalkyl, heterocyclo, or may be fused to a second carbocyclic or heterocyclic ring; R^{4-7} may be hydrogen, halogen, alkoxy(1-6), hydroxy, CN, CO_2H , $CO_2alkyl(1-6)$, $CONHalkyl(1-6)$, $CONDialkyl(1-6)$, NR^9R^{10} , or alkyl(1-6) in which alkyl may be substituted by halogen,
 15 alkoxy(1-6), hydroxy, CN, $CO_2alkyl(1-6)$, $CONHalkyl(1-6)$, $CONDialkyl(1-6)$, NR^9R^{10} , SO_2NR_2 ; any two adjacent substituent R^{4-7} may form a 5-7 membered ring containing 0, 1, or 2 heteroatoms; R^{9-10} may be hydrogen, alkyl(1-6), aryl, heteroaryl, $COCF_3$, SO_2CF_3 , cycloalkyl, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, arylsulfonyl, alkylsulfonyl; or NR^9R^{10} may form a 5 or 6 membered ring such as a pyrrolidine, piperidine, morpholine,
 20 or piperazine ring; n may be 0-3.


US patent 5,175,151 discloses compounds of structure 2 as antiviral and antihypertensive agents,



wherein R^1 may be alkyl, alkenylhalide, $(CH_2)_nCO_2R$, or $(CH_2)_nNR^6R^7$ where R^6 and R^7
 25 may be hydrogen or alkyl; R^2 may include alkyl, alkoxy, aryloxy, aryl, aralkyl, halogen, acyloxy, amino optionally substituted, hydroxy, CH_2OH , CO_2H , alkylesters, CF_3 ; R^3


may be alkyl, aralkyl, optionally substituted aryl, alkylheteroaryl, $(CH_2)_nCO_2R$, or $(CH_2)_nOH$; R^5 may be hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroalkyl, or alkoxy. The patent actually claims a significantly narrower scope which does not include 3-carboxamides.

- 5 PCT publication WO 97/14682 describes compounds useful as gonadotropin-releasing hormone antagonists according to the generic structure 3




- wherein, R^1 is a substituent according to the formula XNR^5R^6 where X is an alkylene or alkyl group optionally substituted by halogen, R^5 is an aralkyl substituent, and R^6 is an alkyl substituent; R^2 is an acylaminoaryl substituent; R^3 is a halogenaralkyl substituent; and R^4 is a carbonyl group optionally esterified or amidated for which $NHCH_2Ph$ is specified in the disclosure.
- 10

- US patent 5,328,887 discloses compounds for the use in a thermal transfer process which would include substructure 4 wherein R may be an optionally substituted hydrocarbon group; R' may be an optionally substituted carboxamide; and the quinoline ring may be optionally substituted.
- 15




PCT publication, WO 98/23608 discloses compounds of structure 5



wherein, among the others, U may include $-(CH_2)_n-$, $-(CH_2)_n-[O,S,N]-(CH_2)_n-$, or linker groups attached through nitrogen; R may be a nitrogen containing heteroaryl substituent or guanidine substituent; R' may be hydrogen, alkyl(1-4), or phenylalkyl(1-4); W is a C₁₋₃ alkyl in which the alkyl group is substituted by either one or two non-hydrogen substituents which may include a halogen substituted phenyl; and Y may be various carboxylate derivatives, sulfonate derivatives, phosphate derivatives and heterocycles, however, may not be hydrogen.


PCT publication, WO99/32450 discloses compounds of structure 6



which are useful as antiviral agents.

SUMMARY OF THE INVENTION

The present invention provides a compound of formula I



15

Wherein R¹ is C₁₋₇ alkyl, optionally substituted by hydroxy or NR⁴R⁵; R² is C₁₋₇ alkyl substituted by hydroxy or NR⁴R⁵; R³ is H, F or C₁₋₇ alkoxy; R⁴ and R⁵ together with N are a 5- or 6-membered heterocyclic moiety having 1-3 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur in which sulfur may be substituted by one (1) or two (2) oxygen atoms; and pharmaceutically acceptable salts thereof.

In another aspect, the present invention also provides:

A pharmaceutical composition which comprises a pharmaceutically acceptable carrier and an effective amount of a compound of formula I,

a method of treating and preventing herpesviral infections in a mammal, including human, and

a use of a compound of formula I to prepare a medicament for treating and preventing herpesviral infections in a mammal, including human.

5

DETAILED DESCRIPTION OF THE INVENTION

For the purpose of the present invention, the carbon atom content of various hydrocarbon-containing moieties is indicated by a prefix designating the minimum and maximum number of carbon atoms in the moiety, i.e., the prefix C_{i-j} indicates a moiety of the integer "i" to the integer "j" carbon atoms, inclusive. Thus, for example, (C₁₋₃)alkyl refers to alkyl of one to three carbon atoms, inclusive, or methyl, ethyl, propyl and isopropyl, straight and branched forms thereof.

The compounds of the present invention are named according to the IUPAC or CAS nomenclature system.

15 The term "C₁₋₇", "C₁₋₅" and "C₁₋₄" alkyl refer to an alkyl group having one to seven, one to five and one to four carbon atoms respectively such as, for example, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl and their isomeric forms thereof.

A 5- or 6-membered heterocyclic moiety includes thiadiazolyl, thiazolyl, 1,1-dioxo-thiazolyl, 1-oxo-thiazolyl, benzothiazolyl, pyridinyl, imidazolyl, indolyl, pyrrolyl, morpholinyl, thiophenyl and 2-oxo-oxazolyl.

20 Compounds of the present invention may be in a form of pharmaceutically acceptable salts.

"Pharmaceutically acceptable salts" refers to those salts which possess the biological effectiveness and properties of the parent compound and which are not biologically or otherwise undesirable. Mammal refers to human and animals.


25 The following Charts A-D describe the preparation of the compounds of formula I of the present invention. All of the starting materials are prepared by procedures described in these charts, by procedures well known to one of ordinary skill in organic chemistry or can be obtained commercially. All of the final compounds of the present invention are prepared by procedures described in these charts or by procedures analogous thereto, which would be well known to one of ordinary skill in organic

30

chemistry. All of the variables used in the charts are as defined below or as in the claims.

In CHART A, structure A-1, ethyl 4-hydroxy-6-iodo-3-quinolinecarboxylate, is prepared by heating 4-iodoaniline with diethyl ethoxymethylene malonate, first at about 150°C, then in refluxing diphenyl ether. Aminolysis of compound A-1 with 4-chlorobenzylamine at about 160°C provides amide A-2. Palladium and copper mediated coupling of A-2 with propargyl alcohol leads to compound A-3. Alkylation of the pyridone nitrogen is accomplished with potassium carbonate and an optionally substituted alkylhalide, affording the compound of structure A-4. Hydrogenation of the alkyne with hydrogen gas, using platinum catalyst, provides the compound of structure A-5.

CHART A



In CHART B, structure B-1, 2-fluoro-5-iodobenzoic acid, is prepared by carbonation of the anion of 4-fluoroiodobenzene. Reaction of B-1 with carbonyldiimidazole, followed by treatment of the resulting acyl imidazolide with ethyl trimethylsilyl malonate and subsequent decarboxylation, affords β -ketoester B-2. The ketoester is converted to quinolinones B-3 by sequential treatment with triethyl orthoformate, an amine, and potassium tert-butoxide. Aminolysis of the ester is accomplished with 4-chlorobenzylamine, giving the compound of structure B-4. Coupling of propargyl alcohol is effected using palladium and copper catalysis, leading

5

The reaction scheme illustrates the synthesis of compounds B-5 and B-6 from B-1. The process begins with B-1 (2-iodo-4-fluorobenzoic acid), which is converted to B-2 (ethyl 2-iodo-4-fluorobenzoylcarbamate). B-2 then undergoes cyclization to form B-3 (ethyl 2-iodo-4-fluoro-1,2,3,4-tetrahydroquinolin-6(1H)-carboxylate). B-3 is further modified to B-4 (ethyl 2-iodo-4-fluoro-1,2,3,4-tetrahydroquinolin-6(1H)-carboxylate, 4-chlorobenzamide derivative). B-4 is then converted to B-5 (ethyl 2-iodo-4-fluoro-1,2,3,4-tetrahydroquinolin-6(1H)-carboxylate, 4-chlorobenzamide derivative, 3-hydroxypropyl derivative). Finally, B-5 is converted to B-6 (ethyl 2-iodo-4-fluoro-1,2,3,4-tetrahydroquinolin-6(1H)-carboxylate, 4-chlorobenzamide derivative, 3-hydroxypropyl derivative, 3-iodopropyl derivative).

B-1 \rightarrow **B-2** \rightarrow **B-3** \rightarrow **B-4**


B-4 \rightarrow **B-5** \rightarrow **B-6**

- 7 -

Chemical reaction scheme showing the synthesis of compound 6 from compound 1:

1 (4-nitro-2,2,4-trimethylphenylacetic acid) \rightarrow 2 (4-nitro-2,2,4-trimethylphenylmethanol) \rightarrow 3 (4-amino-2,2,4-trimethylphenylmethanol benzoate) \rightarrow 4 (4-amino-2,2,4-trimethylphenylmethanol benzoate) \rightarrow 5 (4-amino-2,2,4-trimethylphenylmethanol benzoate) \rightarrow 6 (4-amino-2,2,4-trimethylphenylmethanol benzoate).

CHART D



In CHART E, the corresponding compound of Formula D-4 (see CHART D) where NR₂ is thiomorpholine is reacted with *meta*-chloroperbenzoic acid to afford the sulfoxide (E-1) and the sulfone (E-2).

CHART E





Chart F illustrates a specific method for the preparation of the compound of formula I of the present invention. In CHART F, *N*-methylaniline F-1 is prepared according to a published procedure (Miocque, M; Vierfond, J. M. *Bull. Soc. Chim. Fr.* **1970**, 1901-1907). Compound F-1 is then reacted with ethyl ethoxymethylenemalonate to afford F-2, which is heated in a mixture of Eaton's reagent to afford quinolone F-3. Saponification of F-3 under alkaline conditions affords carboxylic acid F-4. Subsequent coupling of F-4 with 4-chlorobenzylamine mediated by CDI provides F-5. The method is further detailed in Example 3b of the present invention.

CHART F



It will be apparent to those skilled in the art that the described synthetic procedures are merely representative in nature and that alternative synthetic processes are known to one of ordinary skill in organic chemistry.

The examples of the present invention are:

- 5 (a) *N*-(4-Chlorobenzyl)-6-(3-hydroxy-1,1-dimethylpropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- (b) *N*-(4-Chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- (c) *N*-(4-Chlorobenzyl)-1-(2-hydroxyethyl)-6-(3-hydroxypropyl)-4-oxo-1,4-
10 dihydro-3-quinolinecarboxamide;
- (d) *N*-(4-Chlorobenzyl)-6-(3-hydroxypropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- (e) 1-(tert-Butyl)-*N*-(4-chlorobenzyl)-6-(3-hydroxypropyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- 15 (f) *N*-(4-Chlorobenzyl)-6-[(1,1-dioxo-1',4-thiazinan-4-yl)methyl]-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- (g) *N*-(4-Chlorobenzyl)-1-methyl-4-oxo-6-[(1-oxo-1',4-thiazinan-4-yl)methyl]-1,4-dihydro-3-quinolinecarboxamide;
- (h) *N*-(4-chlorobenzyl)-8-fluoro-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-
20 dihydro-3-quinolinecarboxamide; or a pharmaceutically acceptable salt thereof.
- The preferred compounds of the present invention are:
- (a) *N*-(4-Chlorobenzyl)-6-(3-hydroxy-1,1-dimethylpropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- (b) *N*-(4-Chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-
25 3-quinolinecarboxamide;
- (c) *N*-(4-Chlorobenzyl)-1-(2-hydroxyethyl)-6-(3-hydroxypropyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- (d) *N*-(4-Chlorobenzyl)-6-(3-hydroxypropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- 30 (e) *N*-(4-Chlorobenzyl)-6-[(1,1-dioxo-1',4-thiazinan-4-yl)methyl]-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- (f) *N*-(4-Chlorobenzyl)-1-methyl-4-oxo-6-[(1-oxo-1',4-thiazinan-4-yl)methyl]-1,4-dihydro-3-quinolinecarboxamide;

(g) *N*-(4-chlorobenzyl)-8-fluoro-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide; or a pharmaceutically acceptable salt thereof.

The more preferred compounds of the present invention are:

(a) *N*-(4-Chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(b) *N*-(4-Chlorobenzyl)-6-(3-hydroxypropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(c) *N*-(4-Chlorobenzyl)-6-[(1,1-dioxo-1',4-thiazinan-4-yl)methyl]-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(d) *N*-(4-Chlorobenzyl)-1-methyl-4-oxo-6-[(1-oxo-1',4-thiazinan-4-yl)methyl]-1,4-dihydro-3-quinolinecarboxamide;

(e) *N*-(4-chlorobenzyl)-8-fluoro-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide; or a pharmaceutically acceptable salt thereof.

The most preferred compounds of the present invention is *N*-(4-Chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide, or a pharmaceutically acceptable salt thereof.

Also, there is a specific value for the following compounds as synthetic intermediates in the preparation of the preferred compound of formula I:

(a) diethyl 2-([methyl-4-(4-morpholinylmethyl)anilino]methylene)malonate;

(b) ethyl 1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylate; and

(c) 1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylic acid.

The compounds of the present invention and pharmaceutically acceptable salts thereof are useful as antiviral agents. Thus, these compounds are useful to combat viral infections in animals, including man. Specifically, these compounds have anti-viral activity against the herpes virus, cytomegalovirus (CMV). These compounds are also active against other herpes viruses, such as the varicella zoster virus, the Epstein-Barr virus, the herpes simplex virus, and the human herpes virus type 8 (HHV-8).

Also, while many of the compounds of the present invention have shown activity against the CMV polymerase, these compounds may be active against the cytomegalovirus by this or other mechanisms of action. Thus, the description below of

these compounds' activity against the CMV polymerase is not meant to limit the present invention to a specific mechanism of action.

The compounds of the present invention have shown activity in one or more of the assays described below. All of these assays are indicative of a compound's activity and thus of its use as an anti-viral agent.

The HCMV polymerase assay is performed using a scintillation proximity assay (SPA) as described in several references, such as N.D. Cook, et al., Pharmaceutical Manufacturing International, pages 49-53 (1992); K. Takeuchi, Laboratory Practice, September issue (1992); US Patent No. 4,568,649 (1986); which are incorporated by reference herein. Reactions are performed in 96-well plates. The assay is conducted in 100 μ l volume with 5.4 mM HEPES (pH 7.5), 11.7 mM KCl, 4.5 mM $MgCl_2$, 0.36 mg/ml BSA, and 90 nM 3H -dTTP. Assays are run with and without CHAPS, (3-[(3-Cholamidopropyl)-dimethylammonio]-1-propane-sulfonate) at a final concentration of 2 mM. HCMV polymerase is diluted in enzyme dilution buffer containing 50% glycerol, 250 mM NaCl, 10 mM HEPES (pH 7.5), 100 μ g/ml BSA, and 0.01% sodium azide. The HCMV polymerase, which is expressed in recombinant baculovirus-infected SF-9 cells and purified according to literature procedures, is added at 10% (or 10 μ l) of the final reaction volume, i.e., 100 μ l. Compounds are diluted in 50% DMSO and 10 μ l are added to each well. Control wells contain an equivalent concentration of DMSO. Unless noted otherwise, reactions are initiated via the addition of 6 nM biotinylated poly(dA)-oligo(dT) template/primer to reaction mixtures containing the enzyme, substrate, and compounds of interest. Plates are incubated in a 25 C or 37 C H_2O bath and terminated via the addition of 40 μ l/reaction of 0.5 M EDTA (pH 8) per well. Reactions are terminated within the time-frame during which substrate incorporation is linear and varied depending upon the enzyme and conditions used, i.e., 30 min. for HCMV polymerase. Ten μ l of streptavidin-SPA beads (20 mg/ml in PBS/10% glycerol) are added following termination of the reaction. Plates are incubated 10 min. at 37 C, then equilibrated to room temperature, and counted on a Packard Topcount. Linear regressions are performed and IC_{50} 's are calculated using computer software.

A modified version of the above HCMV polymerase assay is performed as described above, but with the following changes: Compounds are diluted in 100% DMSO until final dilution into assay buffer. In the previous assay, compounds are diluted in 50% DMSO. 4.5 mM dithiothreitol (DTT) is added to the polymerase buffer.

Also, a different lot of CMV polymerase is used, which appears to be more active resulting in a more rapid polymerase reaction. Results of the testing of compounds of the present invention in this assay are shown in Tables 1 below. Other viral polymerase assays are performed using procedures similar to those described above.

5 These compounds of the present invention are administered in a pharmaceutical composition containing the compound in combination with a suitable excipient, the composition being useful in combating viral infections. Pharmaceutical compositions containing a compound appropriate for antiviral use are prepared by methods and contain excipients which are well known in the art. A generally recognized compendium
10 of such methods and ingredients is Remington's Pharmaceutical Sciences by E.W. Martin (Mark Publ. Co., 15th Ed., 1975), which is hereby incorporated by reference herein.

 The compounds of the present invention are administered parenterally (for example, by intravenous, intraperitoneal or intramuscular injection), topically, orally, or
15 rectally, depending on whether the preparation is used to treat internal or external viral infections.

 For internal infections, the compositions are administered orally or parenterally at dose levels, calculated as the free base, of about 0.1 to 300 mg/kg, preferably 1.0 to 30 mg/kg of mammal body weight, and are used in man in a unit dosage form, administered
20 one to four times daily in the amount of 1 to 1000 mg per unit dose.

 For parenteral administration or for administration as drops, as for eye infections, the compounds are presented in aqueous solution in a concentration of from about 0.1 to 10%, more preferably about 0.1 to 7%. The solution may contain other ingredients, such as emulsifiers, antioxidants or buffers.

25 The exact regimen for administration of the compounds and compositions disclosed herein will necessarily be dependent upon the needs of the individual subject being treated, the type of treatment and, of course, the judgment of the attending practitioner.


TABLE 1

polymerase IC ₅₀ (μM)			
Example No.	HCMV	HSV	VZV
1	0.5	1.8	2.0
2	7.8	--	--
3	0.4	0.74	0.44
4	0.37	--	--
5	0.48	0.32	0.41
6	2.8	--	--
7	2.5	--	--
8	2.6	--	--
9	4.7	--	--
10	1.0	--	--

The term "--" refers to the data are not determined

5 The compounds and their preparation of the present invention will be better understood in connection with the following examples, which are intended as an illustration of and not a limitation upon the scope of the invention.

EXAMPLE 1. N-(4-chlorobenzyl)-6-(3-hydroxy-1,1-dimethylpropyl)-1-methyl-4-oxo-
10 1,4-dihydro-3-quinolinecarboxamide



To a suspension of 3-methyl-3-(4-nitrophenyl)butanoic acid (16.2 g) (J. Amer. Chem. Soc. 1981, 103, 7768-7773. and J. Amer. Chem. Soc. 1948, 370-371.) in THF (70 mL) is added 103 mL of a 1.0 M borane/THF solution. The reaction is stirred at room
15 temperature for 18 hours. The reaction is poured into H₂O (500 mL). The aqueous layer is saturated with potassium carbonate and extracted with dichloromethane (3 x 500 mL). The combined organic layers are dried with MgSO₄, filtered, and concentrated in vacuo

to yield 16.16 g of the alcohol as a yellow solid. The alcohol (9.41 g) is dissolved in THF (100 mL). Sodium hydride (60% oil dispersion; 5.40 g) is added followed by addition of benzyl bromide (16.0 mL). The reaction is stirred at room temperature for 18 hours. The reaction mixture is partitioned between H₂O (400 mL) and dichloromethane (400 mL). The aqueous layer is extracted with dichloromethane (2 x 400 mL). The combined organic layers are washed with brine (400 mL), dried with MgSO₄, filtered, and concentrated in vacuo. The resulting yellow oil is purified by column chromatography to yield 8.86 g (64%) of the benzyl protected compound as a yellow oil. This material (8.51 g) is suspended in 1/1 conc. HCl/ethanol (200 mL) and SnCl₂•2H₂O (19.22 g) is added. The reaction is heated to 70 °C and is stirred for 2 hours. The reaction mixture is cooled to room temperature, and H₂O (400 mL) is added. The aqueous layer is adjusted to pH 12 with ammonium hydroxide and then extracted with dichloromethane (2 x 400 mL). The combined organic layers are dried with MgSO₄, filtered, and concentrated in vacuo. The resulting brown oil is purified by column chromatography (dichloromethane; dichloromethane/methanol, 98/2) to yield 1.306 g (17%) of the amine as a yellow oil. This material (1.306 g) is combined with diethylethoxymethylenemalonate (0.98 mL) and heated to 120 °C for 2 hours. The reaction mixture is purified by column chromatography (dichloromethane; dichloromethane/methanol, 98/2) to yield 2.044 g (96%) of the malonate intermediate as a yellow oil. This material (2.007 g) is dissolved in diphenyl ether (10 mL) and heated to reflux with removal of ethanol via a Dean-Stark trap for 30 min. The reaction is cooled to room temperature and heptane and ethyl acetate are added. The reaction mixture is allowed to stand at room temperature overnight. The resulting precipitate is filtered off and triturated with ethyl acetate to yield the 0.572 g (32%) of the ester as an off-white solid. The ester (0.517 g) and 4-chlorobenzylamine (1.60 mL) are combined and heated to 190 °C for 1 hour. The reaction is cooled to room temperature and ethyl acetate and heptane are added. The mixture is allowed to stand in the freezer for 3 d. The resulting precipitate is filtered off and recrystallized from ethyl acetate/heptane to yield 0.479 g (75%) of the amide as an off-white solid. To a solution of the amide (0.565 g) in DMF (5 mL) is added potassium carbonate (0.473 g) followed by iodomethane (0.21 mL). The reaction is heated to 90 °C and stirred for 18 hours. The reaction mixture is cooled to room temperature and concentrated in vacuo. The resulting residue is purified by column chromatography (dichloromethane; dichloromethane/methanol 98/2).

Fractions homogeneous by TLC are combined and concentrated in vacuo to yield a pale yellow solid which is recrystallized from ethyl acetate/heptane to yield 0.471 g (81%) of the N-methyl pyridone as a white solid. A solution of the pyridone (0.350 g) in ethanol (50 mL) is hydrogenated over Pd Black (35 mg) for 45 min. The reaction mixture is
5 filtered through a Celite pad and concentrated in vacuo. The resulting yellow solid is recrystallized from ethyl acetate to yield 0.225 g (78%) of the title compound as a pale yellow solid.

Physical characteristics are as follows:

Mp 135-138 °C;

10 ^1H NMR (300 MHz, DMSO- d_6) δ 10.47, 8.85, 8.26, 7.92, 7.79, 7.42-7.35, 4.57, 4.27, 4.04, 3.20, 1.90, 1.35;

^{13}C NMR (75 MHz, DMSO- d_6) δ 176.0, 165.0, 149.0, 146.7, 139.1, 138.5, 131.9, 131.7, 129.6, 128.9, 128.8, 127.8, 126.9, 122.3, 118.0, 110.8, 58.2, 46.6, 41.9, 41.6, 37.0, 29.5;

15 MS (ESI+) for m/z 413 (M+H) $^+$;

IR (drift) 2919, 1916, 1659, 1612, 1555, 1533, 1494, 1431, 1360, 1352, 1085, 813, 806, 797, 682 cm^{-1} .

Anal. Found (corrected for 0.17% H_2O): C, 66.39; H, 6.16; N, 6.65; Cl, 8.39.

20 PREPARATION 1. N-[(4-Chlorophenyl)methyl]-4-hydroxy-6-iodo-3-quinoline-carboxamide.

4-Iodoaniline (8.60 g) and diethyl ethoxymethylenemalonate (7.90 mL) are heated at 130 °C for 1 hour. The reaction is cooled to room temperature and 60 mL diphenyl ether is added. The solution is heated at 250 °C for 1.5 hours with removal of
25 ethanol by a Dean-Stark trap. The reaction is cooled to room temperature and the resulting solid is filtered, washed with hexanes, and dried to yield 11.20 g of ethyl 4-hydroxy-6-iodoquinoline-3-carboxylate. A mixture of this ester (0.58 g) and 4-chlorobenzylamine (4.0 mL) are heated at 180 °C for 1.5 hours. The reaction is cooled and poured into 50 mL diethyl ether. The resulting solid is filtered, triturated in ethyl
30 acetate, and filtered again to give the desired product (0.50 g).

Physical characteristics are as follows:

Mp 297-299 °C;

¹H NMR (300 MHz, DMSO-d₆) 12.71, 10.27, 8.76, 8.50, 8.02, 7.50, 7.38, 7.33, 4.52;

IR (mull) 3151, 3078, 3039, 1631, 1610, 1572, 1563, 1545, 1527, 1512, 1491, 1433, 1351, 1303, 799 cm⁻¹;

5 MS (ES) 438.9 (M+H), 460.9 (M+Na), 436.9 (M-H). Anal. Found: C, 46.61; H, 2.81; N, 6.34; Cl, 8.19.

PREPARATION 2. Methyl 3-{[(4-chlorobenzyl)amino]carbonyl}-4-hydroxy-6-quinolinecarboxylate

10 A solution of N-(4-chlorobenzyl)-4-hydroxy-6-iodo-3-quinolinecarboxamide from Preparation No. 1 (30.0 g), Et₃N (19.1 mL), MeOH (110.6 mL), Pd(OAc)₂ (431 mg), and 1,3-bis (diphenylphosphino) propane (791.9 mg) in 375 mL anhydrous DMF is stirred at room temperature until everything dissolves. CO(g) is slowly bubbled through for 2 days and the reaction is maintained at 70 °C. The reaction is cooled to room
15 temperature. The product is precipitated by adding 160 mL 1N HCl to the reaction mixture. An orange solid precipitates and is collected. The solid is triturated with EtOAc, filtered, and washed with CH₂Cl₂ to afford 23.8 g (93%) of the title compound as an off-white solid.

Physical characteristics are as follows:

20 Mp 290-292 °C;

¹H NMR (300 MHz, DMSO) δ 12.96, 10.26, 8.83, 8.25, 7.80, 7.39, 4.57, 3.9;

IR (drift) 3222, 1724, 1646, 1619, 1574, 1544, 1512, 1489, 1404, 1359, 1288, 1277, 1242, 1210, 738 cm⁻¹;

HRMS (FAB) Found 371.0794. Anal. Found: C, 61.54; H, 3.88; N, 7.51.

25

PREPARATION 3. N-(4-Chlorobenzyl)-4-hydroxy-6-(hydroxymethyl)-3-quinolinecarboxamide.

In a flame-dried 1L 3-necked roundbottom, methyl 3-{[(4-chlorobenzyl)amino]carbonyl}-4-hydroxy-6-quinolinecarboxylate from Preparation No. 2 (3.0 g) is dissolved
30 in 700 mL distilled THF. The suspension is heated to 67 °C to solubilize the starting material. The reaction is allowed to cool to room temperature and then cooled in an ice bath to 10 °C. Lithium aluminum hydride (552.2 mg) is added in one portion. The reaction is stirred at 25 °C and monitored by mass spectroscopy for conversion to desired

product. The reaction is quenched by adding 2 mL H₂O, 2 mL 15% NaOH, and 2 mL H₂O to the reaction mixture. The reaction mixture is filtered to remove the aluminum salt that had precipitated. The filtrate is condensed to obtain a yellow-green residue. The residue is adsorbed onto silica and chromatographed eluting with 2% MeOH in CH₂Cl₂ (1L), 3% MeOH in CH₂Cl₂ (2L), 4% MeOH in CH₂Cl₂ (2L), 5% MeOH in CH₂Cl₂ (1L), 6% MeOH in CH₂Cl₂ (1L), and 7% MeOH in CH₂Cl₂ (2L). The desired product elutes with 4-7% MeOH in CH₂Cl₂. Fractions homogenous by TLC are condensed to yield 1.85 g (67%) of the title compound as yellow crystals.


Physical characteristics are as follows:

10 Mp 288-289 °C;

¹H NMR (300 MHz, DMSO) δ 12.71, 10.48, 8.74, 8.21, 7.71, 7.66, 7.39, 5.38, 4.63, 4.56;

MS (ESI) 343.3 (M+H)⁺, 341.3 (M-H)⁻.

15 EXAMPLE 2. N-(4-Chlorobenzyl)-6-(hydroxymethyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide



A solution of N-(4-chlorobenzyl)-4-hydroxy-6-(hydroxymethyl)-3-quinolinecarboxamide from Preparation No. 3 (300 mg.), K₂CO₃ (485.1 mg), and CH₃I (0.11 mL) in 4 mL anhydrous DMF is heated at 90 °C for 3 hours. The reaction is cooled to room temperature and diluted with H₂O to dissolve any salts and precipitate the product. The crude product is adsorbed onto silica and chromatographed eluting with 3% MeOH in CH₂Cl₂. Fractions homogenous by TLC are combined and condensed to afford 154.2 mg (49%) of the title compound as a white solid.

Physical characteristics are as follows:


Mp 168-170 °C;

¹H NMR (300 MHz, DMSO) δ 10.45, 8.87, 8.30, 7.80, 7.38, 5.42, 4.66, 4.57, 4.02;

30 MS (ESI) 357.2 (M+H)⁺, 355.3 (M-H)⁻.

Anal. Found: C, 63.73; H, 4.62; N, 7.70.

EXAMPLE 3a. N-(4-Chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide.



A solution of N-(4-chlorobenzyl)-6-(hydroxymethyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide from the product of Example 2 (140 mg), collidine (0.061 mL), and DMAP (8.1 mg) in 6.7 mL anhydrous DMF is cooled to 0 °C. Methanesulfonyl chloride (0.12 mL) is added dropwise. The reaction is stirred at room temperature for approx. 2-3 hrs. Morpholine (0.34 mL) is added. The product is precipitated by addition of H₂O. The crude product is adsorbed onto silica and chromatographed eluting with 2% MeOH in CH₂Cl₂. Fractions homogenous by TLC are combined and condensed to afford 70.1 mg (42%) of the title compound as a white solid.

Physical characteristics are as follows:

Mp 188-190 °C;

¹H NMR (300 MHz, DMSO) δ 10.43, 8.87, 8.25, 7.81, 7.38, 4.56, 4.02, 3.63, 3.58, 2.38;

IR (drift) 1654, 1606, 1568, 1554, 1501, 1364, 1348, 1342, 1316, 1134, 1111, 1005, 825, 808, 800 cm⁻¹;

MS (ESI) 426.3 (M+H)⁺, 424.2 (M-H)⁻. Anal. Found: C, 64.61; H, 5.54; N, 9.73.

PREPARATION 4. N-(4-Chlorobenzyl)-4-hydroxy-6-(3-hydroxy-1-propynyl)-3-quinolinecarboxamide.

To a mixture of N-(4-chlorobenzyl)-4-hydroxy-6-iodo-3-quinolinecarboxamide from Preparation No. 1 (0.494 g) in Et₂NH (12.9 mL) is added CuI (10.8 mg) and (Ph₃P)₂PdCl₂ (39.7 mg). DMF (2 mL) is added to solubilize the reactants. To this solution is added propargyl alcohol (0.066 mL) and the reaction is stirred at room temperature for 2 days. The reaction mixture is concentrated to remove Et₂NH. The resulting residue is partitioned between CH₂Cl₂ (3X) and H₂O. A brown solid precipitated from the CH₂Cl₂ layer is filtered and collected to obtain pure product as

indicated by NMR. The organic layers are combined, dried over Na_2SO_4 , and concentrated to obtain a brown residue. The residue is placed under high vac to remove residual DMF. The residue is adsorbed onto silica and chromatographed eluting with 2% MeOH in CH_2Cl_2 and 3% MeOH in CH_2Cl_2 . Fractions homogenous by TLC are combined, condensed and recrystallized with EtOAc/hexanes to obtain a creme solid. The two crops yielded 325.4 mg (79%) of the desired product as a tan solid.

Physical characteristics are as follows:

MP 248-250 C;

^1H NMR (300 MHz, DMSO) 12.85, 10.31, 8.78, 8.22, 7.78, 7.70, 7.38, 5.39, 4.55, 4.33;

IR (drift) 3161, 3073, 3003, 2960, 2914, 1656, 1614, 1557, 1517, 1487, 1299, 1014, 1006, 826, 805 cm^{-1} ;

MS (ESI) 367.0 ($\text{M}+\text{H}^+$), 365.1 ($\text{M}-\text{H}^-$). Anal. Found: C, 65.23; H, 4.24; N, 7.60.

15 PREPARATION 5. N-(4-Chlorobenzyl)-1-(2-hydroxyethyl)-6-(3-hydroxy-1-propynyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide.

To a flask containing N-(4-chlorobenzyl)-4-hydroxy-6-(3-hydroxy-1-propynyl)-3-quinolinecarboxamide from Preparation No. 4 (0.37 g) is added potassium carbonate (2.75 g) and bromoethanol (0.71 mL). The flask is tightly capped and heated to 100 °C.

20 After 4 hours the reaction is cooled to room temperature and partitioned between dichloromethane containing methanol and water. The organic layer is washed with two additional portions of water, brine, dried and concentrated under reduced pressure. The residue is adsorbed onto silica and chromatographed on silica eluting with 2% to 10% methanol in dichloromethane to afford 0.09 g of the title compound as a white solid.


25 Physical characteristics are as follows:

^1H NMR (300 MHz, $\text{DMSO}-d_6$) 10.2, 8.8, 8.3, 7.9, 7.8, 7.4, 5.4, 5.0, 4.5, 4.3, 3.7 ppm;

MS (ESI) m/z 433 ($\text{M}+\text{Na}^+$).

30 EXAMPLE 3b

Alternatively, the compound of EXAMPLE 3a can be prepared as follows:



PREPARATION OF F-2 Diethyl 2-[[methyl-4-(4-morpholinylmethyl)anilino]methylene]malonate;

5 A three liter three-neck flask is charged with F-1 (Miocque, M; Vierfond, J. M. *Bull. Soc. Chim. Fr.* **1970**, 1901-1907), N-(4-methylaminobenzyl)-morpholine (957 g) and diethyl ethoxymethylenemalonate (938 ml) rinsing with ethanol as needed. The solution is heated slowly to about 108° C with atmospheric distillation of ethanol by-product. Distillation is continued until < 5 % morpholine starting material remained as
 10 determined by TLC (silica gel GF; 5 % methanol/methylene chloride). The solution is cooled to room temperature and the residual ethanol is removed by vacuum distillation and toluene azeotrope. High vacuum drying afforded 1778 g (102 % yield) of the title compound F-2 as yellow oil.

¹H NMR (400 MHz CDCl₃) ppm 7.76 (s, 1H), 7.20 (m, 4H), 4.15 (m, 4H), 3.70 (m,
 15 4H), 3.50 (s, 2H), 3.33 (s, 3H), 2.45 (m, 4H) and 1.30 (m, 6H).

¹³C NMR (100 MHz CDCl₃) ppm 167.19, 167.03, 148.72, 145.90, 135.63, 130.12, 121.69, 98.23, 67.00, 62.66, 60.89, 60.27, 53.60, 40.26, 14.41, and 14.15.

PREPARATION OF F-3 Ethyl 1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylate.

A 12 liter three-neck flask is charged with phosphorus pentoxide (294 g) and methanesulfonic acid (1792 ml). The mixture is heated to 50° C with stirring to dissolve the phosphorus pentoxide. The acid solution is cooled to room temperature and a solution of F-2 (780.5 g) in toluene (780.5 ml) is added over 2 hours maintaining the
 25 temperature at < 60° C. After 1 hr at 48-59° C the reaction is complete as determined by TLC (silica gel GF; 5 % methanol/ methylene chloride). The solution is cooled to 0-5°

C and water (780 ml) is added slowly at $< 36^{\circ}\text{C}$. The pH of the slurry is adjusted to 9.5 to 10.5 by addition of 10 M aqueous sodium hydroxide (3365 ml) at $< 36^{\circ}\text{C}$. Water (9330 ml) and methylene chloride (6900 ml) are then added and the product is extracted into the organic layer. The aqueous layer is extracted with methylene chloride (3500 ml). The combined organic layers are washed with water (4000 ml). Removal of the solvent by distillation provided 665 g (97 % yield) of the title compound F-3 as pale yellow crystals.

^1H NMR (400 MHz CDCl_3) ppm 8.42 (m, 2H), 7.40 (m, 1H), 7.39 (m, 1H), 4.38 (q, 2H), 3.88 (s, 3H), 3.68 (m, 4H), 3.62 (s, 2H), 2.47 (m, 4H), and 1.42 (t, 3H).

^{13}C NMR (100 MHz CDCl_3) ppm 174.36, 165.85, 149.47, 138.98, 135.38, 133.58, 128.70, 127.96, 115.80, 111.01, 66.97, 62.53, 60.84, 53.54, 41.35, and 14.44.

PREPARATION OF F-4 1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylic acid.

A 22 liter three-neck flask is charged with F-3 (755 g), ethanol (3776 ml), water (3776 ml), and 50 % sodium hydroxide (831 ml). With stirring under nitrogen the mixture is heated to reflux and maintained until the F-3 is consumed as determined by TLC (silica gel GF; 15 % methanol in methylene chloride, uv). After 1 hr at reflux the reaction is complete. The ethanol is removed by vacuum distillation and the aqueous layer is extracted with methyl t-butyl ether (7550 ml). The pH of the aqueous layer is then lowered to 3.8-4.2 using 6 M hydrochloric acid (2700 ml). The resulting colorless slurry is cooled to 0 to 3°C with stirring. The title compound F-4 is filtered on a coarse sintered glass funnel and washed with 0 to 3°C water (375 ml). The product is dried at 40°C in the vacuum over with nitrogen sweep until $< 0.3\%$ water remained. The yield of F-4 as colorless crystals is 599 g (86.6 % yield). The residual water is 0.245 wt %.

^1H NMR (400 MHz D_2O) ppm 8.70 (s, 1H), 8.39 (s, 1H), 8.11 (m, 1H), 8.02 (m, 1H), 4.83 (s, 3H), 4.66 (s, 2H), 4.05 (m, 4H), and 3.49 (m, 4H).

^{13}C NMR (100 MHz D_2O) ppm 178.06, 168.84, 150.95, 141.26, 137.44, 129.16, 127.42, 125.39, 119.65, 107.47, 64.09, 59.89, 51.85, and 42.64.


PREPARATION OF F-5 N-(4-Chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide.

A 22 liter three-neck flask is charged with compound F-4, (827 g), 1,1'-carbonyldiimidazole (CDI) (443 g), and dimethyl formamide (4960 ml). With stirring
5 under nitrogen the mixture is heated slowly to 60 to 70° C. The solids gradually dissolved as carbon dioxide is steadily evolved. After adding two additional 22 g portions of CDI over about 4 hr, the starting material is consumed as determined by TLC (silica gel GF; 6 % methanol in methylene chloride; uv). As the product of Preparation of F-4 reacted a tan solution is produced, which eventually became a slurry as the acyl
10 imidazolide crystallized. To the slurry is added 4-chlorobenzylamine (333 ml) over 11 min with evolution of heat (64 to 75° C). Dissolution of the imidazolide is followed by crystallization of the product. Stirring is continued at 64 to 80° C until complete as determined by TLC (silica gel GF; 6 % methanol in methylene chloride; uv). An additional 16.7 ml of 4-chlorobenzylamine are added to complete the reaction. On
15 completion, the slurry is cooled to 25° C and 0-5° C water is added (4600 ml). The mixture is cooled to 0-3° C and filtered on a coarse sintered glass funnel. The cake is washed with 0-3° C water (2550 ml) and dried in the vacuum oven at 40°C with a nitrogen sweep. The yield of crude title compound as pale yellow crystals is 947 g (81.3 % yield). The purity by HPLC assay is 97.4 area %. The crude title compound is
20 recrystallized from hot ethanol as needed.

¹H NMR (400 MHz CDCl₃) ppm 10.46 (br s, 1H), 8.77 (s, 1H), 8.41 (s, 1H), 7.80 (m, 1H), 7.49 (m, 1H), 7.30 (m, 4H), 4.64 (d, 2H), 3.95 (s, 3H), 3.70 (m, 4H), 3.64 (s, 2H), and 2.45 (m, 4H).

¹³C NMR (100 MHz CDCl₃) ppm 176.62, 165.07, 148.20, 139.08, 137.40, 135.64,
25 133.94, 132.74, 128.92, 128.60, 127.44, 127.15, 115.99, 111.71, 66.90, 62.50, 53.51, 42.52, and 41.47.

EXAMPLE 4. N-(4-Chlorobenzyl)-1-(2-hydroxyethyl)-6-(3-hydroxypropyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide.



To a solution of N-(4-chlorobenzyl)-1-(2-hydroxyethyl)-6-(3-hydroxy-1-propynyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide from Preparation No. 5 (0.20 g) in a small amount of THF:methanol is added platinum oxide (0.01 g). The mixture is placed under an atmosphere of hydrogen. After 2 hours, the mixture is filtered through Celite with THF:methanol washes. The filtrate is concentrated under reduced pressure. The residue is adsorbed onto silica and chromatographed on silica eluting with 4% to 16% methanol in dichloromethane. The product-containing fractions are concentrated under reduced pressure to afford 0.14 g of the title compound as a white solid.

Physical characteristics are as follows:

^1H NMR (300 MHz, DMSO- d_6) 10.5, 8.7, 8.1, 7.8, 7.7, 7.4, 5.0, 4.5, 3.7, 3.4, 2.7, 1.7;

MS (ESI) m/z 415 ($M+H^+$).

PREPARATION 6. N-(4-Chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide.


A suspension of 6.90 g of N-(4-chlorobenzyl)-4-hydroxy-6-(3-hydroxy-1-propynyl)-3-quinolinecarboxamide from Preparation No. 4, 10.4 g of potassium carbonate, and 2.3 mL of methyl iodide in 40 mL of DMF is stirred at 90°C for 4 hours, then cooled and diluted with 350 mL of water. The resulting solid is filtered, washed well with water, and dried under vacuum. Flash chromatography of the solid on silica using 3-5% methanol in dichloromethane provides 6.02 g of the title compound as a solid.

Physical properties as follows:

^1H NMR ($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ 4.03, 4.45, 4.6, 7.3, 7.6, 7.8, 8.5, 8.8 ppm;

HRMS 381.1006

EXAMPLE 5. N-(4-Chlorobenzyl)-6-(3-hydroxypropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide



A mixture of 0.50 g of N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide from Preparation No. 6 and 50 mg of 5% platinum on carbon catalyst in 20 mL of 1:1 THF-methanol is stirred under 1 atm hydrogen for 3 hours, then filtered through diatomaceous earth. The filtrate is concentrated under reduced pressure and the residual solid flash chromatographed on silica gel using 4-5% methanol in dichloromethane to afford 0.45 g of the title compound as a yellow solid. Further purification is achieved by recrystallization of the solid from 15 mL of acetonitrile.

Physical properties as follows:

^1H NMR ($\text{CDCl}_3 + \text{CD}_3\text{OD}$) δ 1.9, 2.9, 3.6, 4.0, 4.6, 7.3, 7.5, 7.7, 8.3, 8.8 ppm;

HRMS 385.1310.

15

PREPARATION 7. 2-Fluoro-5-iodobenzoic acid.

To an argon-covered, stirred solution of 16.8 mL of diisopropylethylamine in 200 mL of THF, cooled at -78°C , is added dropwise 67 mL of a 1.6 M solution of butyllithium in hexane. The solution is allowed to warm to 0°C and then re-cooled to -78°C . To this solution is added dropwise 11.5 mL of 4-fluoroiodobenzene in 10 mL of THF. The solution is stirred for 90 min at -78°C , then cannulated rapidly onto a Dry Ice-ether slurry. The mixture is allowed to warm to room temperature, then extracted with 300 mL of 0.3 M NaOH. The aqueous phase is chilled in ice and acidified with 40 mL of 6N HCl. The precipitate is extracted with two portions of ether, and the organic phase dried (MgSO_4) and concentrated under reduced pressure. Recrystallization of the residue with ethyl acetate-hexane provides 19.57 g of the title compound as white needles. A second crop of 3.78 g is obtained by recrystallization of the mother liquor residue.

25

Physical properties as follows:

^1H NMR (CDCl_3) δ 6.97, 7.88, 8.33 ppm. Anal found: C, 31.57; H, 1.59.

30

PREPARATION 8. Ethyl 3-(2-fluoro-5-iodophenyl)-3-oxopropanoate.

To a stirred solution of 5.32 g of 2-fluoro-5-iodobenzoic acid from Preparation No. 7 in 20 mL of THF, under argon, is added 3.9 g of carbonyldiimidazole. In a
5 separate flask, 2.8 mL of chlorotrimethylsilane is added to a mixture of 3.74 g of potassium ethyl malonate in 20 mL of acetonitrile. The mixture is stirred under argon for 18 h, then cooled to 0°C for the dropwise addition of 6.6 mL of DBU. The mixture is stirred for 3 h at 0°C, then the solution of acyl imidazolide prepared above is added via cannula. After 2 hours, the mixture is partitioned between ether and excess dilute
10 HCl, and the organic phase is washed with dilute HCl and brine and dried (MgSO₄). Removal of the solvent under reduced pressure left a colorless oil, which is flash chromatographed on silica using 10% ethyl acetate in hexane to provide 5.07 g of the title compound as dense pinkish prisms.

Physical properties as follows:

15 ¹H NMR (CDCl₃) δ 1.34, 4.27, 5.82, 6.89, 7.7, 8.2 ppm;
IR 1624, 1485, 1419, 1245, 1193, 1070, 1028, 813 cm⁻¹.

PREPARATION 9. Ethyl 1-(tert-butyl)-6-iodo-4-oxo-1,4-dihydro-3-quinolinecarboxylate.

20 A solution of 2.36 g of ethyl 3-(2-fluoro-5-iodophenyl)-3-oxopropanoate from Preparation No. 8, 2.0 mL of triethyl orthoformate, and 15 mL of acetic anhydride is refluxed under argon for 2 hours, then the solvents are distilled off under reduced pressure. To the residual oil is added 10 mL of dry tert-butanol and 0.74 mL of tert-butylamine, and the solution is stirred at 80°C for 2 hours. Potassium tert-butoxide (0.87
25 g) is then added, and stirring continued at 80°C under argon for 18 hours. The mixture is then cooled and partitioned between dilute HCl and chloroform-methanol. The organic phase is dried (MgSO₄) and concentrated under reduced pressure. Flash chromatography of the residue on silica using 2-4% methanol in dichloromethane provides 1.32 g of the title compound as an off-white solid.

30 Physical properties as follows:

¹H NMR (CDCl₃) δ 1.42, 1.87, 4.4, 7.7, 7.9, 8.9 ppm;
HRMS 400.0414. Anal. Found: C, 48.05; H, 4.50; N, 3.52.

PREPARATION 10. 1-(tert-Butyl)-N-(4-chlorobenzyl)-6-iodo-4-oxo-1,4-dihydro-3-quinolinecarboxamide.

A slurry of 1.11 g of ethyl 1-(tert-butyl)-6-iodo-4-oxo-1,4-dihydro-3-quinolinecarboxylate from Preparation No. 9 in 2.0 g of 4-chlorobenzylamine is heated
5 under argon at 160°C for 18 hours, then cooled to room temperature and triturated with 1N HCl. The solid is filtered, washed well with water, and dried under vacuum. Flash chromatography using 20% ethyl acetate in dichloromethane provides 1.22 g of the title compound as a white solid.

Physical properties as follows:

10 ¹H NMR (CDCl₃) δ 1.89, 4.6, 7.3, 7.7, 7.9, 8.9, 9.22, 10.4 ppm;

IR 1664, 1536, 1468, 1342, 1180 cm⁻¹.

Anal. Found: C, 51.27; H, 4.19; N, 5.62.

PREPARATION 11. 1-(tert-Butyl)-N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide
15

To a stirred slurry of 1.15 g of 1-(tert-butyl)-N-(4-chlorobenzyl)-6-iodo-4-oxo-1,4-dihydro-3-quinolinecarboxamide from Preparation No. 10, 156 mg of copper (I) iodide, and 66 mg of dichlorobis(triphenylphosphine)palladium (II) in 23 mL of diethylamine, under argon, is added 0.16 mL of propargyl alcohol. The mixture is stirred
20 for 18 hour at room temperature, then concentrated under reduced pressure. The residue is partitioned between water and chloroform-methanol, and the organic phase dried (MgSO₄) and concentrated under reduced pressure. Flash chromatography of the residue on silica using 2-4% methanol in dichloromethane affords 977 mg of tan solid.

Recrystallization from ethanol provides 850 mg of the title compound as a beige solid.


25 Physical properties as follows:

¹H NMR (CDCl₃) δ 1.92, 4.47, 4.6, 7.3, 7.7, 8.0, 8.5, 9.19, 10.5 ppm;

HRMS 423.1466.

Anal. Found: C, 67.74; H, 5.53; N, 6.61.

30 EXAMPLE 6. 1-(tert-Butyl)-N-(4-chlorobenzyl)-6-(3-hydroxypropyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide.



A mixture of 303 mg of 1-(tert-butyl)-N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide from Preparation No. 11 and 15 mg of platinum oxide in 10 mL of 1:1 THF-methanol is stirred under 1 atm of hydrogen gas for 3 h, then filtered through diatomaceous earth and concentrated under reduced pressure. The mixture was purified by flash chromatography on silica using 2-3% methanol in dichloromethane to afford 294 mg of the title compound.


Physical properties as follows:

^1H NMR (CDCl_3) δ 1.89, 1.9, 2.9, 3.7, 4.6, 7.3, 7.5, 7.9, 8.4, 9.21, 10.6 ppm;

IR 1658, 1596, 1548, 1484, 1349, 1184, 810, 731 cm^{-1} ;

HRMS 427.1762

EXAMPLE 7. *N*-(4-Chlorobenzyl)-1-methyl-4-oxo-6-(4-thiomorpholinyl-methyl)-1,4-dihydro-3-quinolinecarboxamide.




Methanesulfonyl chloride (0.193 mL) is added to a solution of *N*-(4-chlorobenzyl)-6-(hydroxymethyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide (357 mg) from Example No. 2, DMAP (20 mg), and 2,4,6-collidine (0.33 mL) in anhydrous DMF (20 mL). The mixture is stirred at room temperature for 2 h, and poured into water (60 mL). The resulting precipitate is filtered, washed with water (20 mL), and recrystallized from acetonitrile to afford 0.328 g (74%) of the title compound as a white solid.

Physical characteristics are as follows:

Mp 215-219 $^{\circ}\text{C}$; ^1H NMR ($\text{DMSO}-d_6$) δ 10.42, 8.87, 8.23, 7.80, 7.42-7.34, 4.56, 4.02, 3.66, 2.62; ^{13}C NMR ($\text{DMSO}-d_6$) δ 175.4, 164.4, 148.7, 139.0, 138.7, 135.3, 133.7, 131.3, 129.1, 128.3, 126.7, 125.6, 117.6, 110.5, 61.8, 54.3, 41.4, 41.2, 27.2; IR (drift) 2915, 1655, 1605, 1574, 1551, 1502, 1364, 1339, 1316, 1132, 826, 808, 801, 726, 661

cm⁻¹; MS (ESI+) *m/z* 442 (M+H)⁺. Anal. found for C₂₃H₂₄ClN₃O₂S: C, 62.18; H, 5.46; N, 9.47; Cl, 8.40; S, 7.02.


- EXAMPLES 8-9. *N*-(4-Chlorobenzyl)-1-methyl-4-oxo-6-(4-thiomorpholinyl-methyl)-1,4-dihydro-3-quinolinecarboxamide (221 mg) from Example No. 7 is dissolved in dry CH₂Cl₂ (15 mL) and the solution is cooled to 0°C. A solution of mCPBA (258 mg) in dry CH₂Cl₂ (5 mL) is added dropwise, maintaining the temperature of the solution between 0-2°C. The reaction is stirred at 0°C for 2 h and then is allowed to warm to room temperature. The reaction is quenched with saturated NaHSO₃ (2 mL) and stirred for another hour. Dilution with water (10 mL) forms a white precipitate which is filtered. The crude solid is a mixture of the two products which are purified by column chromatography (MeOH/CH₂Cl₂, 1/49 to 3/97) to afford 37 mg (16%) of Example No. 8 and 54 mg (24%) of Example No. 9 as white solids.
- EXAMPLE 8. *N*-(4-chlorobenzyl)-6-[(1,1-dioxo-1',4-thiazinan-4-yl)methyl]-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide.



Physical characteristics are as follows:

- Mp 245-247°C; ¹H NMR (DMSO-*d*₆) δ 10.42, 8.88, 8.26, 7.83, 7.42-7.34, 4.56, 4.03, 3.85, 3.11, 2.90; IR (drift) 1916, 1662, 1610, 1569, 1539, 1498, 1326, 1317, 1289, 1268, 1123, 1107, 805, 796, 661 cm⁻¹. Anal. Found for C₂₃H₂₄ClN₃O₄S: C, 58.05; H, 5.06; N, 8.78; Cl, 7.55.


EXAMPLE 9. *N*-(4-chlorobenzyl)-1-methyl-4-oxo-6-[(1-oxo-1',4-thiazinan-4-yl)methyl]-1,4-dihydro-3-quinolinecarboxamide.



Physical characteristics are as follows:

Mp 249-251°C; ¹H NMR (DMSO-*d*₆) δ 10.42, 8.87, 8.26, 7.82, 7.42-7.34, 4.56, 4.02, 3.74, 2.91-2.83, 2.74-2.62; IR (drift) 2818, 2351, 1974, 1932, 1656, 1604, 1574, 1551, 1500, 1363, 1336, 1054, 1027, 838, 809 cm⁻¹. Anal. Found for C₂₃H₂₄ClN₃O₃S: C, 60.09; H, 5.30; N, 9.03; Cl, 7.90.

EXAMPLE 10. *N*-(4-chlorobenzyl)-8-fluoro-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide




To a solution of *N*-(4-chlorobenzyl)-8-fluoro-4-hydroxy-6-(4-morpholinylmethyl)-3-quinolinecarboxamide, (can be prepared according to the procedure described in PCT patent application, WO99/32450) (147 mg) in anhydrous DMF (3 mL) is added K₂CO₃ (71.2 mg) followed by CH₃I (0.026 mL). The reaction mixture is stirred at room temperature for 30 min, then poured into H₂O (40 mL) to precipitate the product. The solid is collected, adsorbed onto silica, and chromatographed (1% MeOH in CH₂Cl₂ (2L)). Fractions homogenous by TLC are combined and concentrated to afford 55.3 mg of the title compound as a white solid.

Physical characteristics are as follows:

Mp 187-189 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.28, 8.77, 8.09, 7.65, 7.38, 4.56, 4.17, 3.58, 2.38; IR (drift) 1659, 1603, 1578, 1556, 1550, 1543, 1499, 1366, 1348, 1278, 1134, 1117, 865, 809, 797 cm⁻¹; HRMS (FAB) calcd for C₂₃H₂₃ClFN₃O₃+H₁ 444.1490, found 444.1484.

CLAIMS

1. A compound of formula I



I

5 Wherein

R^1 is C_{1-7} alkyl, optionally substituted by hydroxy or NR^4R^5 ;

R^2 is C_{1-7} alkyl substituted by hydroxy or NR^4R^5 ;

R^3 is H, F or C_{1-7} alkoxy;

10 R^4 and R^5 together with N are a 5- or 6-membered heterocyclic moiety having 1-3 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur in which sulfur may be substituted by one (1) or two (2) oxygen atoms; and a pharmaceutically acceptable salt thereof.

2. A compound of claim 1 wherein R^1 is C_{1-4} alkyl, optionally substituted by hydroxy;
15 R^2 is C_{1-5} alkyl substituted by hydroxy or morpholine; and R^3 is H or F.

3. A compound of claim 1 which is

- (a) *N*-(4-chlorobenzyl)-6-(3-hydroxy-1,1-dimethylpropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- 20 (b) *N*-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- (c) *N*-(4-chlorobenzyl)-1-(2-hydroxyethyl)-6-(3-hydroxypropyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- (d) *N*-(4-chlorobenzyl)-6-(3-hydroxypropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- 25 (e) 1-(tert-butyl)-*N*-(4-chlorobenzyl)-6-(3-hydroxypropyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide;
- (f) *N*-(4-chlorobenzyl)-6-[(1,1-dioxo-1',4-thiazinan-4-yl)methyl]-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(g) *N*-(4-chlorobenzyl)-1-methyl-4-oxo-6-[(1-oxo-1',4-thiazinan-4-yl)methyl]-1,4-dihydro-3-quinolinecarboxamide;

(h) *N*-(4-chlorobenzyl)-8-fluoro-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide; or a pharmaceutically acceptable salt thereof.

5

4. A compound of claim 1 which is

(a) *N*-(4-chlorobenzyl)-6-(3-hydroxy-1,1-dimethylpropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(b) *N*-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-
10 dihydro-3-quinolinecarboxamide;

(c) *N*-(4-chlorobenzyl)-1-(2-hydroxyethyl)-6-(3-hydroxypropyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(d) *N*-(4-chlorobenzyl)-6-(3-hydroxypropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(e) *N*-(4-chlorobenzyl)-6-[(1,1-dioxo-1',4-thiazinan-4-yl)methyl]-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(f) *N*-(4-chlorobenzyl)-1-methyl-4-oxo-6-[(1-oxo-1',4-thiazinan-4-yl)methyl]-1,4-dihydro-3-quinolinecarboxamide;

(g) *N*-(4-chlorobenzyl)-8-fluoro-1-methyl-6-(4-morpholinylmethyl)-4-oxo-
20 1,4-dihydro-3-quinolinecarboxamide; or a pharmaceutically acceptable salt thereof.

5. A compound of claim 1 which is

(a) *N*-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(b) *N*-(4-chlorobenzyl)-6-(3-hydroxypropyl)-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(c) *N*-(4-chlorobenzyl)-6-[(1,1-dioxo-1',4-thiazinan-4-yl)methyl]-1-methyl-4-oxo-1,4-dihydro-3-quinolinecarboxamide;

(d) *N*-(4-chlorobenzyl)-1-methyl-4-oxo-6-[(1-oxo-1',4-thiazinan-4-yl)methyl]-1,4-dihydro-3-quinolinecarboxamide;

(e) *N*-(4-chlorobenzyl)-8-fluoro-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide; or a pharmaceutically acceptable salt thereof.

6. A compound of claim 1 which is
N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxamide, or a pharmaceutically acceptable salt thereof.
- 5 7. A compound of formula I for use in treating infections from herpesviruses which comprises administering to a patient in need thereof an effective amount of a compound of formula I as shown in claim 1.
8. The use of claim 7 wherein said herpesviruses is herpes simplex virus
10 types 1, herpes simplex virus types 2, varicella zoster virus, cytomegalovirus, Epstein-Barr virus, human herpes viruses 6, human herpes viruses 7 or human herpes viruses 8.
9. The use of claim 7 wherein said herpesviruses is herpes simplex virus
15 types 1, herpes simplex virus types 2, varicella zoster virus, cytomegalovirus, Epstein-Barr virus, human herpes viruses 7 or human herpes viruses 8.
10. The use of claim 7 wherein said herpesviruses is human cytomegalovirus.
- 20 11. The use of claim 7 wherein the effective amount of a compound of claim 1 is administered orally, parenterally or topically.
12. The use of claim 7 wherein the effective amount of a compound of claim 1 is in an amount of from about 0.1 to about 300 mg/kg of body weight.
25
13. The use of claim 7 wherein the effective amount of a compound of claim 1 is in an amount of from about 1 to about 30 mg/kg of body weight.
14. A pharmaceutical composition which comprises an effective amount of a
30 compound of claim 1 and a pharmaceutically acceptable carrier.

15. A use of the compound of formula I for the preparation of a medicament for inhibiting a viral DNA polymerase comprising contacting the polymerase with an effective inhibitory amount of a compound of formula I as shown in claim 1.
- 5 16. Compounds useful as synthetic intermediates in the preparation of the compound of claim 6 which are
- (a) diethyl 2-([methyl-4-(4-morpholinylmethyl)anilino]methylene)malonate;
- (b) ethyl 1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylate; and
- 10 (c) 1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylic acid.
17. A process for the preparation of the compound of claim 6 comprising the steps of
- a. heating a mixture of diethyl ethoxymethylenemalonate and N-(4-methyl-aminobenzyl)morpholine to provide diethyl 2-([methyl-4-(4-morpholinylmethyl)anilino]methylene)malonate,
- 15 b. adding a solution of diethyl 2-([methyl-4-(4-morpholinylmethyl)anilino]methylene)malonate in toluene to a mixture of phosphorus pentoxide and methanesulfonic acid to provide ethyl 1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylate,
- 20 c. converting ethyl 1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylate to 1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylic acid,
- d. stirring 1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylic acid with carbonyldiimidazole and dimethyl formamide, and
- 25 e. reacting 4-chlorobenzylamine with the mixture of step d.

INTERNATIONAL SEARCH REPORT

Int'l. Application No
PCT/US 99/27959

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D215/56 C07D417/06 A61K31/47 A61P31/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	WO 99 32450 A (PHARMACIA& UPJOHN COMPANY) 1 July 1999 (1999-07-01) cited in the application page 222 -page 252; claims 1-17	1-17
A	WO 97 04779 A (CHIROSCIENCE LIMITED) 13 February 1997 (1997-02-13) page 3, line 28 -page 5, line 8 page 30 -page 36; claims 1-27	1-17
A	WO 92 04328 A (SCHERING CORPORATION) 19 March 1992 (1992-03-19) page 1 -page 5, line 10	1-17
A	FR 2 593 810 A (SANOFI SOCIÉTÉ ANONYME) 7 August 1987 (1987-08-07) page 1 -page 3	1-17
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

20 April 2000

Date of mailing of the international search report

02/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/27959

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 175 151 A (ADRIANO AFONSO ET AL.) 29 December 1992 (1992-12-29) cited in the application page 1 -page 6	1-17
A	US 4 959 363 A (MARK P. WENTLAND) 25 September 1990 (1990-09-25) page 1 -page 3, line 50	1-17
A	US 4 786 644 A (EDWARD J. GLAMKOWSKI ET AL.) 22 November 1988 (1988-11-22) the whole document	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. J. Application No

PCT/US 99/27959

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9932450 A	01-07-1999	AU 1606399 A	12-07-1999
WO 9704779 A	13-02-1997	AU 696390 B	10-09-1998
		AU 6626396 A	26-02-1997
		CA 2225552 A	13-02-1997
		EP 0841929 A	20-05-1998
		JP 11513021 T	09-11-1999
		US 5891878 A	06-04-1999
		AU 695132 B	06-08-1998
		AU 6626696 A	26-02-1997
		CA 2225555 A	13-02-1997
		EP 0841927 A	20-05-1998
		WO 9704775 A	13-02-1997
		JP 11510156 T	07-09-1999
		US 5753666 A	19-05-1998
WO 9204328 A	19-03-1992	AT 161252 T	15-01-1998
		AU 8448491 A	30-03-1992
		AU 8665991 A	30-03-1992
		CA 2091172 A,C	08-03-1992
		CA 2091173 A	08-03-1992
		DE 69128481 D	29-01-1998
		DE 69128481 T	09-04-1998
		EP 0550464 A	14-07-1993
		EP 0549729 A	07-07-1993
		JP 6502845 T	31-03-1994
		JP 7110853 B	29-11-1995
		US 5412104 A	02-05-1995
		WO 9204327 A	19-03-1992
		US 5942522 A	24-08-1999
		US 5378694 A	03-01-1995
FR 2593810 A	07-08-1987	AT 52082 T	15-05-1990
		AU 587102 B	03-08-1989
		AU 6791687 A	30-07-1987
		CA 1310324 A	17-11-1992
		DK 35387 A	25-07-1987
		EP 0234995 A	02-09-1987
		GR 3000440 T	28-06-1991
		JP 62185074 A	13-08-1987
		NZ 218918 A	26-04-1989
		PT 84172 A,B	01-02-1987
		US 4835163 A	30-05-1989
		ZA 8700348 A	26-08-1987
US 5175151 A	29-12-1992	NONE	
US 4959363 A	25-09-1990	NONE	
US 4786644 A	22-11-1988	DK 658688 A	28-05-1989
		EP 0317991 A	31-05-1989
		JP 2138260 A	28-05-1990
		PT 89094 A,B	01-12-1988
		US 4966906 A	30-10-1990
		US 4952588 A	28-08-1990

